


# On the Rainfall and Temperature Forecast Skill for a Tropical Andean Mountainous Area in Northern South America Using Different Operational Weather Forecast Strategies: Role of the Diurnal Cycle of Rainfall on the Success of Data Assimilation.


Mauricio Zapata<sup>(1,2)</sup>, Gisel Guzmán-Echavarría<sup>(1)</sup>, Carlos D. Hoyos<sup>(1,2)</sup>, Juan C. Hernández<sup>(1)</sup>, Lina I. Ceballos<sup>(1)</sup>, Manuela Guarín<sup>(1)</sup> 1. Sistema de Alerta temprana de Medellín y el Valle de Aburrá 2. Universidad Nacional de Colombia, Sede Medellín.

## Introduction

As a part of a regional risk management strategy, we designed two different operational weather forecast strategies for a complex terrain region in the Andes mountain range in northern South America. Both strategies, together, generate a total of eleven different forecasts every day, using the initial and boundary conditions from the GFS.



Figures (a) and (b) show the geographical context of the Aburrá valley. Figures (c), (d) and (e), show a north, centre and south cross-section of the valley. Figure (f), (g) and (h) show the diurnal cycle of rainfall (Blue) and temperature (Red), obtained from the SIATA's rain gauge and AWS network.



Diurnal cycle of precipitation inside of the study region, using the meteorological radar dBz retrievals.

## Model setup

### SYNOPSIS

Medium-range forecast  
Forecast lead time: 120 hours  
Runs per day: 3 runs  
Input data: GFS 0.5 12 UTC  
Domain: Triple nested domain  
Micro-physics: Purdue Lin scheme (PAR02), Eta Ferrier scheme (PAR05) and Thompson Scheme (PAR08)

### RDFS

Short-range forecast  
Forecast lead time: 30 hours  
Runs per day: 8 runs (4 with and 4 without data assimilation)  
Input data: GFS 00, 06, 12 and 18 UTC and Doppler radar C-band data for data assimilation  
Domain: Double nested domain  
Micro-physics: Thompson Scheme (PAR08)

Both configurations: Short-range (dashed white line) and Medium-range (solid white line), share the continental domain as can be seen in figure (a). Figure (b) shows the inner domain boundaries for both configurations. The shaded circle corresponds radar coverage. Figure (c) shows a cross-section at 6.2 north degrees. The magenta star in (b) and (c) shows the location of the radar.





Figure (a) shows the operational forecast timeline for both configurations. Figure (b) shows the daily timeline of historical configurations (short and mid-range forecast). Short-range forecast or RDFS starts every day at eight different times, every six hours with (DA) and without data assimilation (CTRL). The mid-range forecast or SYNOPSIS starts every day with the 12 UTC boundary conditions.

## Rainfall forecast verification


To determine the spatial differences between the rainfall forecast (SYNOPSIS and RDFS) and the rainfall observations (radar), we quantify the mean absolute error (MAE) for the cumulative rainfall during the early morning, morning, afternoon and night.




Spatial distribution of MAE. Each column corresponds to a forecast configuration. The first three columns corresponds to SYNOPSIS, RDFS CTRL and DA respectively for every initial hour (00, 06, 12 and 18 UTC). The rows represent the early morning, morning, afternoon and night.

## Temperature forecast verification

There is a systematic error in the surface temperature forecast due to the difference between the real and model topography. This BIAS is corrected every day in the operational post-processing using a Model Output Statistics (MOS) scheme.




Monthly diurnal cycle of temperature differences between forecast and observations with and without MOS



Elevation differences vs. temperature differences Crosses (points) indicate temperature differences before (after) MOS.

## Surface Energy budget



Observed, simulated and differences of surface radiation (land use = Urban)

## Diurnal cycle representation






Figure shows the spatial distribution of hit rate at night for the Aburrá Valley. The north zone (Barbosa) shows the worst skill, mainly in the data assimilation runs.



Mean absolute Error (MAE), per hour in the forecast lead time on rdfs configuración. In the Center of Aburrá Valley. From left to the right 00, 06, 12, and 18 UTC.

## Overall conclusions and future work

- At the inner domain, the best configuration is RDFS 12 UTC control run, but inside of Aburrá Valley, there are notable differences in the skill due to local differences in the representation of the rainfall diurnal cycle.
- Due to similarities in the performance between runs of SYNOPSIS configuration, regarding RDFS ones, we conclude that skill of rainfall forecast is highly dependent of atmospheric input data (also data assimilation) more than the choice of microphysics schema.
- The problems on the temperature representation are remarkable, due to biases in shortwave radiation and turbulent fluxes. However, temperature forecasts are accurate after a statistical correction (MOS).

## Acknowledgments

This work was supported by Area Metropolitana de Medellín y del Valle de Aburrá, Municipio de Medellín, Grupo EPM, and ISAGEN under the contract CCT504 of 2019.